



WIC* Counselors Children

Yogurt, milk, and calcium-fortified soymilk contain calcium that children need!

CHILDREN NEED CALCIUM

- Most bone mass development happens in childhood.^{1,2}
- Calcium helps build strong bones.³
- Nearly 1/3 of WIC children (aged 3-5) don't get enough calcium.⁴
- Preschool children get much less calcium after age two.⁵

OTHER BENEFITS

- Yogurt, milk, and calcium-fortified soymilk have high-quality protein kids need for muscles and growth.
- Calcium-fortified soymilk is naturally lactose free.
- Yogurt may be a good option for a lactose intolerant child.
 - Yogurt often has less lactose than milk.⁷ (Greek yogurt has even less.)
- Full fat yogurt, and whole milk, have fat children 1-2 need for growth and development.⁸⁻¹⁰

YOGURT, MILK, AND CALCIUM-FORTIFIED SOYMILK PROVIDE CALCIUM AND MORE

- A 1-cup serving of yogurt or milk is usually a good source of calcium.
- A 1-cup serving of calcium-fortified unsweetened soymilk is a good non-dairy option.
- Yogurt, milk, and calcium-fortified soymilk often have vitamin D added.
- 83% of WIC children (3-5) don't get enough vitamin D.⁶
Vitamin D and calcium work together to help build strong bones.³

HELPFUL HINTS

- Babies under 1 shouldn't drink milk or soymilk.
- Kids 1-3 need 2 to 2-1/2 servings of dairy per day.¹¹
- Kids 4-5 need 2-1/2 to 3 servings of dairy per day.¹¹
- Kids under 2 need full fat dairy. After 2, give low fat or non fat dairy.
- Use milk or calcium-fortified unsweetened soymilk in cereal or oatmeal.
Cook with yogurt. Get recipes at: danonewave.com/WIC

References

1. NIH News in Health. (2010). Keeping Bones Strong and Healthy: Let's Talk About Osteoporosis. [online] Available at: <http://newsinhealth.nih.gov/2010/February/feature1.htm> [Accessed 13 Dec. 2017]. NIH Osteoporosis and Related Bone Diseases. National Resource Center.
2. NIH Osteoporosis and Related Bone Diseases National Resource Center. (2015). Osteoporosis: Peak Bone Mass in Women. [online] Available at: https://www.bones.nih.gov/sites/bones/files/bone_mass.pdf [Accessed 13 Dec. 2017].
3. National Institutes of Health. Office of Dietary Supplements. (2017). - Calcium. [online] Available at: <https://ods.od.nih.gov/factsheets/Calcium-HealthProfessional/> [Accessed 13 Dec. 2017].
4. Data analysis NHANES (National Health and Nutrition Examination Survey) 2009-2010.
5. U.S. Department of Agriculture, Agriculture Research Service. Data tables: results from USDA's 1994-96 Continuing Survey on Food Intakes by Individuals and 1994-96 Diet and Knowledge Survey.
6. Institute of Medicine. (2006). WIC Food Packages: Time for a Change. Washington, DC: The National Academies Press. [online] Available at: <https://doi.org/10.17226/11280> [Accessed 13 Dec. 2017].
7. Lomer, et al. Aliment Pharmacol Ther. 2008;27(2):93-103.
8. Hardy and Kleinman. J Pediatr. 1994;125(5 Pt 2):S69-77.
9. Kleinman's Pediatric Nutrition Handbook, 5th Ed. American Academy of Pediatrics, 2004, p. 126.
10. Allen RE, Myers AL. Nutrition in Toddlers. Am Fam Physician. 2006 Nov 1;74(9):1527-32.
11. 2015-2020 Dietary Guidelines. (2015). A Closer Look at Current Intakes and Recommended Shifts. [online] Available at: <https://health.gov/dietaryguidelines/2015/guidelines/chapter-2/a-closer-look-at-current-intakes-and-recommended-shifts/> [Accessed 13 Dec. 2017].

*WIC is a registered service mark of the U.S. Department of Agriculture for USDA's Special Supplemental Nutrition Program for Women, Infants and Children.